

Trögerite**(UO₂)₃(AsO₄)₂·12H₂O(?)**

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Tetragonal or monoclinic. *Point Group:* 4/m 2/m 2/m or 2/m. As thin crystals, tabular on {001}, to 1 mm.

Physical Properties: *Cleavage:* On {001}, perfect; on {100}, good; on {011}, probable. Hardness = 2–3 D(meas.) = 3.3 D(calc.) = [5.13] Reversible H₂O content. May fluoresce lemon-yellow under UV. Radioactive.

Optical Properties: Transparent. *Color:* Lemon-yellow. *Luster:* Vitreous, pearly on {001} cleavage.

Optical Class: Uniaxial (-) or biaxial (-), probably anomalous, zoned due to differences in H₂O content. *Pleochroism:* O = Y = Z = lemon-yellow; E = X = nearly colorless. *Orientation:* X = c; Z ∧ a = 12°–14°. *Dispersion:* r > v, moderate. ω = 1.624–1.627 ε = 1.580–1.582 α = 1.584–1.600 β = 1.620–1.630 γ = 1.623–1.630 2V(meas.) = 0°–40°

Cell Data: *Space Group:* n.d. a = 6.99 c = 8.48 Z = 8, or *Space Group:* n.d. a = 7.15 b = 6.98 c = 11.2 β = ~95° Z = 1

X-ray Powder Pattern: Synthetic.

8.59 (10), 3.79 (9), 3.30 (8), 5.50 (7), 4.35 (7), 2.70 (7), 2.19 (7)

Chemistry:

	(1)	(2)	(3)
UO ₃	63.76	65.17	65.80
P ₂ O ₅		1.20	
As ₂ O ₅	19.64	22.08	17.62
Bi ₂ O ₃		8.79	
PbO		1.56	
MgO		0.25	
CaO		0.65	
H ₂ O	14.81		16.58
Total	98.21	99.70	100.00

(1) Weisser Hirsch mine, Germany. (2) Do.; by electron microprobe, after drying at 210 °C; corresponds to (UO₂)_{2.72}Bi_{0.18}Ca_{0.11}Pb_{0.07}Mg_{0.06}[(AsO₄)_{1.84}(PO₄)_{0.16}]_{Σ=2.00}.

(3) (UO₂)₃(AsO₄)₂·12H₂O.

Mineral Group: Autunite group.

Occurrence: A rare mineral formed in the oxidation zone of some uranium deposits.

Association: Walpurgite, uranospinite, uranosphathite, asselbornite, zeunerite, uranosphaerite, erythrite, cobaltian wad (Weisser Hirsch mine, Germany); realgar, orpiment, scorodite, mansfieldite, sodium uranospinite, arseniosiderite, metatorbernite, metazeunerite, uranophane, arsenopyrite, pyrite, galena (Bota-Burum deposit, Kazakhstan).

Distribution: From the Walpurgis vein, in the Weisser Hirsch mine, and at the Daniel mine, Neustädtel-Schneeberg, Saxony, Germany. From Marbrue, near Dalbeattie, Kirkcudbrightshire, Scotland. In the Rabéjac, Riviéral, and Mas-d'Alary uranium deposits, near Lodève, Hérault, France. At the Bota-Burum uranium deposit, 15 km southwest of Alakol Lake, Chu-Ili Mountains, southwestern Balkhash district, Kazakhstan. From the Bald Mountain district, Lawrence Co., South Dakota, USA.

Name: To honor Mining Foreman R. Tröger, Schneeberg, Germany, who found the first specimens.

Type Material: Mining Academy, Freiberg, Germany, 21800; Harvard University, Cambridge, Massachusetts, USA, 106035.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 966–967. (2) Frondel, C. (1958) Systematic mineralogy of uranium and thorium. U.S. Geol. Sur. Bull. 1064, 187–191. (3) Weiss, A., F. Taborszky, K. Hartl, and E. Tröger (1957) Zur Kenntnis des Uranminerals Trögerite. Zeits. Naturforsch., 12b, 356–358.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.